

Complete Summary

GUIDELINE TITLE

Acne.

BIBLIOGRAPHIC SOURCE(S)

Finnish Medical Society Duodecim. Acne. In: EBM Guidelines. Evidence-Based Medicine [Internet]. Helsinki, Finland: Wiley Interscience. John Wiley & Sons; 2007 Jan 4 [Various].

GUIDELINE STATUS

Note: This guideline has been updated. The National Guideline Clearinghouse (NGC) is working to update this summary.

** REGULATORY ALERT **

FDA WARNING/REGULATORY ALERT

Note from the National Guideline Clearinghouse: This guideline references a drug(s) for which important revised regulatory and/or warning information has been released.

- [October 6, 2006, iPLEDGE](#): An update to iPLEDGE that will eliminate one element of the program.
- [August 12, 2005, iPLEDGE](#): Approval of a strengthened risk management program, called iPLEDGE, for Accutane and generic isotretinoin.

COMPLETE SUMMARY CONTENT

** REGULATORY ALERT **

SCOPE

METHODOLOGY - including Rating Scheme and Cost Analysis

RECOMMENDATIONS

EVIDENCE SUPPORTING THE RECOMMENDATIONS

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

CONTRAINDICATIONS

IMPLEMENTATION OF THE GUIDELINE

INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT

CATEGORIES

IDENTIFYING INFORMATION AND AVAILABILITY

DISCLAIMER

SCOPE

DISEASE/CONDITION(S)

Acne, including:

- Comedonic acne (A. comedonicus)
- Common acne (A. vulgaris) or pustular acne
- Cystic acne (A. cystica)
- Acne conglobata
- Acne fulminans

GUIDELINE CATEGORY

Treatment

CLINICAL SPECIALTY

Dermatology
Family Practice
Internal Medicine
Pediatrics

INTENDED USERS

Health Care Providers
Physicians

GUIDELINE OBJECTIVE(S)

Evidence-Based Medicine Guidelines collect, summarize, and update the core clinical knowledge essential in general practice. The guidelines also describe the scientific evidence underlying the given recommendations.

TARGET POPULATION

Patients with acne

INTERVENTIONS AND PRACTICES CONSIDERED

Treatment

1. Skin cleansing with soap or antibacterial detergents
2. Treatment of comedonic acne with retinoic acid cream or solution, adapalen gel, or benzoyl peroxide cream or gel (3 to 10%)
3. Treatment of common acne with local antibiotics (e.g., clindamycin solution), ultraviolet light therapy, and systemic treatment as indicated
4. Systemic antibiotics (tetracycline, erythromycin)
5. Local treatment and light therapy, used simultaneously with systemic treatment

6. Incision and drainage of pus-containing cysts with a large-caliber injection needle or narrow-tipped scalpel
7. Hormonal treatment for women: Cyproterone acetate (an anti-androgen) + oestrogen
8. Treatment of scars by skin abrasion or laser therapy (by a dermatologist or a plastic surgeon)
9. Isotretinoin upon recommendation of a dermatologist
10. Consultation with or referral to a dermatologist

Note: Guideline developers considered several other treatment options. For a list of these, see "Related Resources" in the original guideline document and the "Major Recommendations" field below.

MAJOR OUTCOMES CONSIDERED

- Efficacy of treatment
- Adverse effects of treatment

METHODOLOGY

METHODS USED TO COLLECT/SELECT EVIDENCE

Hand-searches of Published Literature (Primary Sources)
 Hand-searches of Published Literature (Secondary Sources)
 Searches of Electronic Databases

DESCRIPTION OF METHODS USED TO COLLECT/SELECT THE EVIDENCE

The evidence reviewed was collected from the Cochrane database of systematic reviews and the Database of Abstracts of Reviews of Effectiveness (DARE). In addition, the Cochrane Library and medical journals were searched specifically for original publications.

NUMBER OF SOURCE DOCUMENTS

Not stated

METHODS USED TO ASSESS THE QUALITY AND STRENGTH OF THE EVIDENCE

Weighting According to a Rating Scheme (Scheme Given)

RATING SCHEME FOR THE STRENGTH OF THE EVIDENCE

Levels of Evidence

- A. Strong research-based evidence. Multiple relevant, high-quality scientific studies with homogeneous results.
- B. Moderate research-based evidence. At least one relevant, high-quality study or multiple adequate studies.
- C. Limited research-based evidence. At least one adequate scientific study.

D. No research-based evidence. Expert panel evaluation of other information.

METHODS USED TO ANALYZE THE EVIDENCE

Systematic Review

DESCRIPTION OF THE METHODS USED TO ANALYZE THE EVIDENCE

Not stated

METHODS USED TO FORMULATE THE RECOMMENDATIONS

Not stated

RATING SCHEME FOR THE STRENGTH OF THE RECOMMENDATIONS

Not applicable

COST ANALYSIS

A formal cost analysis was not performed and published cost analyses were not reviewed.

METHOD OF GUIDELINE VALIDATION

Peer Review

DESCRIPTION OF METHOD OF GUIDELINE VALIDATION

Not stated

RECOMMENDATIONS

MAJOR RECOMMENDATIONS

Note: This guideline has been updated. The National Guideline Clearinghouse (NGC) is working to update this summary. The recommendations that follow are based on the previous version of the guideline.

The levels of evidence [A-D] supporting the recommendations are defined at the end of the "Major Recommendations" field.

Classification of Acne

- Comedonic acne (A. comedonicus, See Figure 1 in the original guideline document)
 - Plenty of open or obstructed comedos, but scant inflammatory changes

- Common acne (*A. vulgaris*) or pustular acne (See Figures 2 & 3 in the original guideline document)
 - Pustules and comedos
- Cystic acne (*A. Cystica*, See Figures 4 & 5 in the original guideline document)
 - Cystic foci of infection that result in scars
- Acne conglobata
 - Multilobular inflammatory cysts containing volatile pus
 - Therapy-resistant, scar forming
- Acne fulminans
 - An uncommon variant of acne in young men characterized by systemic symptoms (fever, arthralgia, skeletal foci of inflammation)
 - Systemic corticosteroids, not antibiotics, are the drugs of choice.
 - Refer patients with suspected acne fulminans to a dermatologist without delay. The painful disease is not well known, and is often left untreated for a long time.

Treatment

Local Treatment

- Local treatment is usually sufficient for comedonic acne and mild common acne.
- Wash the skin with soap or antibacterial detergents.
- Comedonic acne can be treated with
 - Retinoic acid cream or solution (tretinoin [Purdy, 2005] [A], isotretinoin[[Purdy, 2005] [B]])
 - Adapalen gel (Purdy, 2005) [C]
 - Benzoyl peroxide (3 to 10%) (Purdy, 2005) [A] cream or gel
 - All above drugs can be irritating at first. Use a low concentration of the active drug initially, and advise the patient to wash the drug away after a few hours. The tolerance of the skin increases with time.
- Common acne can be treated with
 - Local antibiotics (e.g., clindamycin solution) (Purdy, 2005) [A]
 - Ultraviolet light therapy (as a course of 15 treatments added to other treatment) for widespread disease
- Consider systemic treatment if the effect of local treatment is unsatisfactory 2 to 3 months from the onset of treatment.

Systemic Treatment

- Antibiotics
 - Tetracycline (Garner et al., 2003) [B] and erythromycin (Purdy, 2005) [A] are equally effective. The usual dose is 250 to 500 mg/day for a few months. Six months' treatment with tetracycline or erythromycin 1 g/day is more effective than a shorter treatment with a smaller dose. Do not use tetracyclines in children below 12 years of age.
 - Local treatment and light therapy can be used simultaneously with systemic treatment.
 - Local treatment is not sufficient in cystic acne and conglobate acne. Use systemic antibiotics or consider referral to a dermatologist. Pus-containing cysts can be drained by incising them with a large-caliber injection needle or narrow-tipped scalpel.

- Hormonal treatment for women
 - Cyproterone acetate (an anti-androgen) + oestrogen for 6 months reduce the excretion of sebaceous glands and alleviate acne.

Acne Scars

- Consider treatment of scars by skin abrasion or laser therapy (Jordan, Cummins, & Burls, West Midlands, 1998; Jordan, Cummins, & Burls, Birmingham, 1998; Health Technology Assessment Database [HTA]-998502, 2001) [D] only after the activity of the disease has totally subsided.
- Scars can be treated either by a dermatologist or a plastic surgeon.

Indications for Specialist Consultation

- Severe forms of acne (A. cystica, conglobata, fulminans)
- If ordinary treatment fails, the dermatologist can consider isotretinoin. However, it has considerable teratogenicity.

Related Resources

See original guideline document for Cochrane reviews and other evidence summaries.

Definitions:

Levels of Evidence

- Strong research-based evidence. Multiple relevant, high-quality scientific studies with homogeneous results.
- Moderate research-based evidence. At least one relevant, high-quality study or multiple adequate studies.
- Limited research-based evidence. At least one adequate scientific study.
- No research-based evidence. Expert panel evaluation of other information.

CLINICAL ALGORITHM(S)

None provided

EVIDENCE SUPPORTING THE RECOMMENDATIONS

REFERENCES SUPPORTING THE RECOMMENDATIONS

[References open in a new window](#)

TYPE OF EVIDENCE SUPPORTING THE RECOMMENDATIONS

Concise summaries of scientific evidence attached to the individual guidelines are the unique feature of the Evidence-Based Medicine Guidelines. The evidence summaries allow the clinician to judge how well-founded the treatment

recommendations are. The type of supporting evidence is identified and graded for select recommendations (see the "Major Recommendations" field).

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

POTENTIAL BENEFITS

Effective treatment of acne

POTENTIAL HARMS

Adverse Effects of Medication

- Retinoic acid cream or solution, adapalene gel, and benzoyl peroxide (3 to 10%) can be irritating at first. The tolerance of the skin increases with time.
- Isotretinoin has considerable teratogenicity

CONTRAINDICATIONS

CONTRAINDICATIONS

Tetracyclines should not be used in children below 12 years of age.

IMPLEMENTATION OF THE GUIDELINE

DESCRIPTION OF IMPLEMENTATION STRATEGY

An implementation strategy was not provided.

INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

IOM CARE NEED

Getting Better

IOM DOMAIN

Effectiveness

IDENTIFYING INFORMATION AND AVAILABILITY

BIBLIOGRAPHIC SOURCE(S)

Finnish Medical Society Duodecim. Acne. In: EBM Guidelines. Evidence-Based Medicine [Internet]. Helsinki, Finland: Wiley Interscience. John Wiley & Sons; 2007 Jan 4 [Various].

ADAPTATION

Not applicable: The guideline was not adapted from another source.

DATE RELEASED

2001 Apr 30 (revised 2007 Jan 4)

GUIDELINE DEVELOPER(S)

Finnish Medical Society Duodecim - Professional Association

SOURCE(S) OF FUNDING

Finnish Medical Society Duodecim

GUIDELINE COMMITTEE

Editorial Team of EBM Guidelines

COMPOSITION OF GROUP THAT AUTHORED THE GUIDELINE

Primary Author: Jorma Lauharanta

FINANCIAL DISCLOSURES/CONFLICTS OF INTEREST

Not stated

GUIDELINE STATUS

Note: This guideline has been updated. The National Guideline Clearinghouse (NGC) is working to update this summary.

GUIDELINE AVAILABILITY

This guideline is included in "EBM Guidelines. Evidence-Based Medicine" available from Duodecim Medical Publications, Ltd, PO Box 713, 00101 Helsinki, Finland; e-mail: info@ebm-guidelines.com; Web site: www.ebm-guidelines.com.

AVAILABILITY OF COMPANION DOCUMENTS

None available

PATIENT RESOURCES

None available

NGC STATUS

This summary was completed by ECRI on December 17, 2002. The information was verified by the guideline developer as of February 7, 2003. This summary was updated by ECRI on March 29, 2004, and again on September 29, 2004. This summary was updated by ECRI on June 19, 2006 following the U.S. Food and Drug Administration (FDA) advisory on Accutane and generic isotretinoin. This NGC summary was updated by ECRI on August 7, 2006. This summary was updated by ECRI on November 14, 2006 following the updated FDA advisory on iPLEDGE. This NGC summary was updated by ECRI on February 27, 2007.

COPYRIGHT STATEMENT

This NGC summary is based on the original guideline, which is subject to the guideline developer's copyright restrictions.

DISCLAIMER

NGC DISCLAIMER

The National Guideline Clearinghouse™ (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at <http://www.guideline.gov/about/inclusion.aspx>.

NGC, AHRQ, and its contractor ECRI make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.

© 1998-2007 National Guideline Clearinghouse

Date Modified: 10/22/2007

